

CANTOR SETS IN THE LOGISTIC FAMILY

W. PATRICK HOOPER

We consider the logistic map $F_\mu(x) = \mu x(1 - x)$ when $\mu > 4$. Define

$$\Lambda = \{x \in [0, 1] : F_\mu^n(x) \in [0, 1] \text{ for all } n \geq 0\}.$$

We will prove the following theorem:

Theorem 1. *If $\mu > 2 + \sqrt{5}$, then Λ is a Cantor set.*

The goal of this note is to give a more explicit proof of this result than what is offered in Devaney's book. The proof that Λ is perfect is different than what we did in class, and basically just is a more detailed account of Devaney's proof.

We will now give an equivalent description of Λ . Define the set

$$A_0 = \{x \in [0, 1] : F_\mu(x) \notin [0, 1]\}.$$

Then A_0 is an open interval:

$$A_0 = \left(\frac{1}{2} - \frac{\sqrt{\mu - 4}}{2\sqrt{\mu}}, \frac{1}{2} + \frac{\sqrt{\mu - 4}}{2\sqrt{\mu}} \right).$$

We can then inductively define A_n for all integers $n \geq 0$ inductively according to the rule that $A_{n+1} = F_\mu^{-1}(A_n)$. Then we have

$$\Lambda = [0, 1] \setminus \bigcup_{n=0}^{\infty} A_n.$$

The reason for taking $\mu > 2 + \sqrt{5}$ is given by the following, which is not hard to prove.

Proposition 2. *If $\mu > 2 + \sqrt{5}$, then $|F'_\mu(x)| > 1$ for all $x \in [0, 1] \setminus A_0$.*

The relevance of the derivative is given by the following, which follows from the fundamental theorem of calculus and was proved in class.

Proposition 3. *If f is a real-valued function which is differentiable and monotone on an interval $J \subset \mathbb{R}$, then*

$$\text{length } f(J) = \int_J |f'(x)| dx.$$

These two statements are enough to prove our Theorem.

Proof of Theorem 1. Suppose that $\mu > 2 + \sqrt{5}$. We will prove that Λ is closed, totally disconnected, and perfect.

1. We will prove that Λ is closed. Observe that A_0 is open. Recall that $A_{n+1} = F_\mu^{-1}(A_n)$. Since F_μ is continuous, $F_\mu^{-1}(U)$ is open for every open set $U \subset \mathbb{R}$. Therefore, we can see by induction that A_n is open for all n . Then $\Lambda = [0, 1] \setminus \bigcup_{n=0}^{\infty} A_n$ must be closed.

2. We will prove that Λ is totally disconnected, that is Λ does not contain any (non-degenerate) intervals. By proposition 2, there is a $\lambda > 1$ so that $|F'_\mu(x)| \geq \lambda$ for all $x \in [0, 1] \setminus A_0$.

Suppose Λ does contain an interval J_0 of positive length. Then, inductively define J_n for all $n \geq 0$ according to the rule that $J_{n+1} = F_\mu(J_n)$. This makes $J_n = F_\mu^n(J_0)$ for all n . Since $J_0 \subset \Lambda$, we know that $J_n \subset \Lambda$ for all n . Then, it must be that each J_n lies in one of the two closed intervals making up $[0, 1] \setminus A_0$. Since F_μ is differentiable and monotone on each of these intervals, we can use Proposition 3 to see that

$$\text{length } J_{n+1} = \int_{J_n} |F'_\mu(x)| dx \geq \int_{J_n} \lambda dx = \lambda \cdot \text{length } J_n.$$

It follows that the length of J_n grows exponentially. But this is impossible since $J_n \subset [0, 1]$ for all n .

3. We will prove that Λ is perfect. It is enough to prove that for all $x_0 \in \Lambda$ and all $\epsilon > 0$ there is a $y \in \Lambda$ so that $x_0 \neq y$ and $|x_0 - y| < \epsilon$.

The set $[0, 1] \setminus A_0$ consists of two closed intervals,

$$I_0 = \left[0, \frac{1}{2} - \frac{\sqrt{\mu-4}}{2\sqrt{\mu}}\right] \quad \text{and} \quad I_1 = \left[\frac{1}{2} + \frac{\sqrt{\mu-4}}{2\sqrt{\mu}}, 1\right].$$

The restrictions of F_μ to I_0 and I_1 ,

$$F_\mu|_{I_0} : I_0 \rightarrow [0, 1] \quad \text{and} \quad F_\mu|_{I_1} : I_1 \rightarrow [0, 1],$$

are homeomorphisms, so they have inverse maps

$$h_0 : [0, 1] \rightarrow I_0 \quad \text{and} \quad h_1 : [0, 1] \rightarrow I_1.$$

Observe that if $x \in \Lambda$ then $h_0(x) \in \Lambda$ and $h_1(x) \in \Lambda$. This is because

$$x = F_\mu(h_0(x)) = F_\mu(h_1(x)).$$

So, the orbits of $h_0(x)$ and $h_1(x)$ follow the orbit of x .

By Proposition 3, we know that the restrictions $F_\mu|_{I_0}$ and $F_\mu|_{I_1}$ expand distance by a factor of $\lambda > 1$. So, h_0 and h_1 each contract distances by a factor of λ . That is, if $a, b \in [0, 1]$, then

$$|h_0(a) - h_0(b)| \leq \frac{1}{\lambda}|a - b| \quad \text{and} \quad |h_1(a) - h_1(b)| \leq \frac{1}{\lambda}|a - b|.$$

Choose any $x_0 \in \Lambda$. Then the orbit $\{x_n = F_\mu^n(x_0) : n \geq 0\}$ is contained in Λ for all $n \geq 0$. Since $\Lambda \subset I_0 \cup I_1$, there is a sequence $\langle s_n \in \{0, 1\} : n \geq 0 \rangle$ so that

$$F_\mu^n(x) \in I_{s_n} \quad \text{for all } n \geq 0.$$

In particular, notice that since $F_\mu(x_n) = x_{n+1}$, we know that $x_n = h_{s_n}(x_{n+1})$, since $x_n \in I_{s_n}$ and the image of x_n under F_μ is x_{n+1} . It then follows that

$$x_0 = h_{s_0} \circ \dots \circ h_{s_{n-2}} \circ h_{s_{n-1}}(x_n).$$

Now pick any $\epsilon > 0$. Then there is an n so that $\frac{1}{\lambda^n} < \epsilon$. Now choose $y' = 0$ or $y' = 1$, so that $y' \neq x_n$. Note that both points lie in Λ , since they map to zero which is fixed by F_μ . Define

$$y = h_{s_0} \circ \dots \circ h_{s_{n-2}} \circ h_{s_{n-1}}(y').$$

Since each map is a homeomorphism, we know that $x_0 \neq y$. Also these maps contract distances by $\frac{1}{\lambda}$. So

$$|x_0 - y| \leq \frac{1}{\lambda^n} |x_n - y'| \leq \frac{1}{\lambda^n} < \epsilon.$$

□

THE CITY COLLEGE OF NEW YORK, NEW YORK, NY, USA 10031
E-mail address: whooper@ccny.cuny.edu