

Math A4500: Midterm 2 Study Guide

Prof. Hooper

Disclaimer. This test is just a recommendation of things to study. You may be asked about things that do not appear here. I recommend ensuring that you can do all the problems in the sections covered in the book.

Definitions. You will be asked to define several terms on the test. The following is a list of terms which might appear.

Cantor set, totally disconnected, perfect, sequence space (or shift space), shift map, itinerary (or symbolic code), topological conjugacy, topologically transitive, sensitive dependence on initial conditions, chaotic, topologically semi-conjugate, C^r -distance, C^r -structurally stable

Theorems. Theorems and results from the book you should be able to prove.

- Propositions 4.4 and 4.6: Local behaviors near a hyperbolic fixed point.
- Propositions 5.1-5.3: Results about the Logistic map when $1 < \mu < 3$.
- Proposition 5.6: If $\mu > 2 + \sqrt{5}$ then Λ is a Cantor set.
- Proposition 6.6: Dynamical facts about the shift map on Σ_2 .
- Theorem 9.8: The local behavior of a map near a hyperbolic fixed point is locally conjugate to its derivative.

Techniques. You should be able to use the various techniques we have developed for understanding particular dynamical systems acting on subsets of the real line. For instance, you should be able to:

- Prove that a subset of the real line is a Cantor set. This set could be constructed or dynamically determined.
- Prove dynamical results about a Shift Space. See Proposition 6.6.
- Prove that the dynamics of f on a set Λ is topologically conjugate to a shift map on a shift space.
- Prove that a system is chaotic.
- Prove two systems are topologically conjugate using fundamental domains. This was done in class, but also see example 9.4.
- Use topological conjugacy to prove dynamical statements about a map. We discussed in class how f and g are topologically conjugate then they have similar dynamical properties. See Theorem 7.5, for instance.
- Prove a system is C^r -structurally stable, or not C^r structurally stable for a fixed r and a fixed map f .

- Sharkovsky's theorem: You should be able to use the techniques from §1.10 to find periodic orbits for a continuous map $f : \mathbb{R} \rightarrow \mathbb{R}$.

Types of problems. There will be between four and six multi-part problems on the midterm. You'll have the full class period (100 minutes) to complete the midterm. Problems of the following forms are likely to appear:

- *Homework:* Problems similar to assigned homework problems.
- *Recall something, then prove something:* Problems that ask you to recall a definition or a major result, and then use it in a proof.
- *Math comprehension:* The problem states a new definition, which may not have been seen before, and asks you to use the definition in basic proofs.
- *Prove a result:* Problems which ask you to prove a result which is proved in the book. See the section titled "Theorems" above.

Practice problems. These are some problems I gave in a midterm last time I taught this course.

1. **Motivation:** Recall that we showed that when $\mu > 2 + \sqrt{5}$ that the logistic map $F_\mu(x) = \mu x(1 - x)$ has an invariant Cantor set. The set was defined by

$$\Lambda = \{x \in [0, 1] : F_\mu^n(x) \in [0, 1] \text{ for all integers } n \geq 0\}.$$

One fact we needed to prove was that Λ is closed. This problem asks you to generalize this proof.

Prove the following statement:

Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be a continuous function, and let $X \subset \mathbb{R}$ be an arbitrary closed set. Prove that

$$\Lambda = \{x \in X : f^n(x) \in X \text{ for all integers } n \geq 0\}$$

is a closed set.

2. (a) (10 points) Suppose $f : \mathbb{R} \rightarrow \mathbb{R}$ is a homeomorphism so that $f(x) > x$ for all $x \in \mathbb{R}$. Prove that $\lim_{n \rightarrow \infty} f^n(x) = +\infty$ for all $x \in \mathbb{R}$.
 - (b) (6 points) Complete the following definition:
Let $f : A \rightarrow A$ and $g : B \rightarrow B$ be two maps with $A, B \subset \mathbb{R}$. We say f and g are *topologically conjugate* if . . .
 - (c) Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be a homeomorphism so that $f(x) > x$ for all $x \in \mathbb{R}$ be as in the prior part, and let $g : \mathbb{R} \rightarrow \mathbb{R}$ be $g(x) = x + 1$. Prove that f and g are topologically conjugate.
(*Hint:* Use part (a) and the similar fact that $\lim_{n \rightarrow \infty} f^{-n}(x) = -\infty$ for all $x \in \mathbb{R}$. This second fact can be assumed without proof.)

Recall the following definitions and use them in the following proofs.

- The C^r -distance between two C^r functions $f, g : \mathbb{R} \rightarrow \mathbb{R}$ is

$$d_r(f, g) = \sup_{x \in \mathbb{R}} \max_{k=0, \dots, r} |f^{(k)}(x) - g^{(k)}(x)|.$$

- A C^r function $g : \mathbb{R} \rightarrow \mathbb{R}$ is C^r -structurally stable if there is an $\epsilon > 0$ so that f and g are topologically conjugate whenever f is a C^r function satisfying $d_r(f, g) < \epsilon$.
- (d) Prove that $g(x) = x + 1$ is C^1 -structurally stable. (*Hint:* You can use part (c), even if you were not successful in writing a proof for that part.)
- (e) Prove that $g(x) = x + 1$ is not C^0 -structurally stable.

3. Suppose $f : \mathbb{R} \rightarrow \mathbb{R}$ is continuous and has the following orbit of period four:

$$f(0) = 2, \quad f^2(0) = 3, \quad f^3(0) = 1, \quad f^4(0) = 0.$$

Define the the intervals $A = [0, 1]$, $B = [1, 2]$ and $C = [2, 3]$.

- (a) Prove that f has a point of least period 13 which stays within $A \cup B \cup C$. Find an explicit sequence of intervals visited by such an orbit.
- (b) Use Sharkovsky's theorem to prove that f has points of all possible periods. (*Remarks:* It is easier to use Sharkovsky's theorem then to prove this fact with the methods used to prove Sharkovsky's theorem, however the application of Sharkovsky's theorem is indirect. You can use the lemmas from the proof of Sharkovsky's theorem without proving the lemmas.)
4. Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be differentiable function with finitely many critical points. Prove that $f^m(x)$ also has only finitely many critical points for every $m \geq 1$.
5. (The Logistic Map) Let $F(x) = \mu x(1 - x)$ with $\mu > 2 + \sqrt{5}$. Let

$$\Lambda = \{x : F_\mu^n(x) \in [0, 1] \text{ for all } n \geq 0\}.$$

Let $p_0 \in (0, \frac{1}{2})$ and $p_1 \in (\frac{1}{2}, 1)$ be the points so that $f(p_0) = f(p_1) = 1$. Define $I_0 = [0, p_0]$ and $I_1 = [p_1, 1]$. Then, $\Lambda \subset I_0 \cup I_1$, and $|F'(x)| > k$ for some $k > 1$ and all $x \in I_0 \cup I_1$.

- (a) (5 points) Define *sensitive dependence on initial conditions*.
- (b) (5 points) Describe how to code the orbit of a point in Λ using a one-sided shift space. That is, describe the map $s : \Lambda \rightarrow \Sigma_2$.
- (c) (5 points) Prove that the coding map s is injective.
- (d) (5 points) Assuming that the coding map s is a topological conjugacy between $F|_\Lambda : \Lambda \rightarrow \Lambda$ and $\sigma : \Sigma_2 \rightarrow \Sigma_2$, prove that F has a dense set of periodic points. (You do not need to prove that σ has a dense set of periodic points.)

6. (Shift spaces) Let $\mathcal{A} = \{0, 1\}$. Define the one-sided shift space

$$\Sigma = \{(s_0 s_1 s_2 \dots) : s_j \in \mathcal{A} \text{ for all integers } j \geq 0\}.$$

The shift space becomes a metric space when equipped with either of the following distance functions:

$$d(\mathbf{s}, \mathbf{t}) = \sum_{i=0}^{\infty} \frac{|s_i - t_i|}{2^i} \quad \text{or} \quad d'(\mathbf{s}, \mathbf{t}) = \begin{cases} 2^{-n} & \text{if } \mathbf{s} \neq \mathbf{t} \text{ and } n = \min\{i : s_i \neq t_i\} \\ 0 & \text{iff } \mathbf{s} = \mathbf{t}. \end{cases}$$

Both d and d' determine the same topology. The shift map $\sigma : \Sigma \rightarrow \Sigma$ is defined by

$$\sigma(s_0 s_1 s_2 \dots) = (s_1 s_2 s_3 \dots).$$

Use the definitions above and either of the metrics to answer the following questions.

- (a) (5 points) Is the map σ a homeomorphism? Why or why not?
- (b) (5 points) Prove that σ -periodic points are dense in Σ .
- (c) (5 points) Define *topologically transitive*.
- (d) (5 points) Prove that the shift map is topologically transitive on Σ .